

Von Hippel-Lindau and Hereditary Pheochromocytoma/Paraganglioma Syndromes: Clinical Features, Genetics, and Surveillance Recommendations in Childhood



Surya P. Rednam¹, Ayelet Erez², Harriet Druker³, Katherine A. Janeway⁴, Junne Kamihara⁴, Wendy K. Kohlmann⁵, Katherine L. Nathanson⁶, Lisa J. States⁷, Gail E. Tomlinson⁸, Anita Villani³, Stephan D. Voss⁹, Joshua D. Schiffman^{5,10}, and Jonathan D. Wasserman¹¹

Abstract

Von Hippel-Lindau disease (vHL) is a hereditary tumor predisposition syndrome that places affected individuals at risk for multiple tumors, which are predominantly benign and generally occur in the central nervous system or abdomen. Although the majority of tumors occur in adults, children and adolescents with the condition develop a significant proportion of vHL manifestations and are vulnerable to delayed tumor detection and their sequelae. Although multiple tumor screening paradigms are currently being utilized for patients with vHL, surveillance should be reassessed as the available relevant clinical information continues to expand. We propose a new vHL screening paradigm similar to existing approaches, with important modifications for some tumor types, placing an emphasis on risks in childhood. This includes advancement in the timing of surveillance initiation and increased frequency of screening evaluations. Another neuroendocrine-related familial condition is the rapidly expanding

hereditary paraganglioma and pheochromocytoma syndrome (HPP). The tumor spectrum for patients with HPP syndrome includes paragangliomas, pheochromocytomas, renal cancer, and gastrointestinal stromal tumors. The majority of patients with HPP syndrome harbor an underlying variant in one of the *SDHx* genes (*SDHA*, *SDHB*, *SDHC*, *SDHD*, *SDHA*, and *SDHAF2*), although other genes also have been described (*MAX* and *TMEM127*). Annual screening for elevated plasma or urine markers along with complete blood count and biennial whole-body MRI accompanied by focal neck MRI is recommended for older children and adults with HPP syndrome to detect tumors early and to decrease morbidity and mortality from HPP-related tumors. *Clin Cancer Res*; 23(12); e68–e75. ©2017 AACR.

See all articles in the online-only CCR Pediatric Oncology Series.

¹Department of Pediatrics, Baylor College of Medicine, Texas Children's Cancer Center, Texas Children's Hospital, Houston, Texas. ²Department of Biological Regulation, Weizmann Institute of Science, Rehovot, Israel. ³Division of Haematology/Oncology, The Hospital for Sick Children, Department of Paediatrics, University of Toronto, Toronto, Ontario, Canada. ⁴Department of Pediatric Oncology, Dana-Farber Cancer Institute and Children's Hospital, Boston, Massachusetts. ⁵Huntsman Cancer Institute, University of Utah, Salt Lake City, Utah. ⁶Department of Medicine, Division of Translational Medicine and Human Genetics, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, Pennsylvania. ⁷Department of Radiology, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania. ⁸Department of Pediatrics, Division of Hematology and Oncology and Greehey Children's Cancer Research Institute, University of Texas Health Science Center at San Antonio, San Antonio, Texas. ⁹Department of Radiology, Children's Hospital, Boston, Massachusetts. ¹⁰Department of Pediatrics, University of Utah, Salt Lake City, Utah. ¹¹Division of Endocrinology, The Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada.

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

S.P. Rednam and A. Erez contributed equally to this article.

Corresponding Author: Jonathan D. Wasserman, The Hospital for Sick Children, 555 University Avenue, Toronto, Ontario, Canada M5G 1X8. Phone: 416-813-7654, ext. 228186; Fax: 416-813-6304; E-mail: jonathan.wasserman@sickkids.ca

doi: 10.1158/1078-0432.CCR-17-0547

©2017 American Association for Cancer Research.

Von Hippel-Lindau Disease

Introduction

Von Hippel-Lindau disease (vHL; OMIM #193300) is a multisystemic tumor predisposition syndrome characterized by benign and malignant tumors, including central nervous system (CNS) and retinal hemangioblastomas, clear cell renal cell carcinoma (RCC), pheochromocytoma (PHEO), pancreatic neuroendocrine tumors (pancreatic NET), endolymphatic sac tumor (ELSI), and epididymal and broad ligament cystadenoma, as well as visceral (renal and pancreatic) cysts (1). Originally described in 1936 in the context of familial retinal angiomatosis by von Hippel (2) and CNS hemangioblastomas by Lindau (3), the incidence of vHL is estimated at approximately one in 36,000 (based on data preceding the advent of molecular genetic testing), and the lifetime penetrance approaches 100% by age 75 (4).

A clinical diagnosis of vHL can be established in one of two scenarios: (i) in an individual with a family history of vHL and the presence of a CNS or retinal hemangioblastoma, PHEO, or RCC; or (ii) in a simplex case (an individual with no family history) with ≥ 2 hemangioblastomas or ≥ 2 visceral tumors or one hemangioblastoma and one visceral tumor (5). Clinically, vHL is subdivided into five subtypes based on tumor spectrum and

Table 1. Clinical phenotypes of vHL are classified into four types

	Type I	Type IB	Type IIA	Type IIB	Type IIC
Clinical manifestations	<ul style="list-style-type: none"> Retinal angioma CNS HB RCC Pancreatic NETs Low risk for PHEO 	<ul style="list-style-type: none"> Retinal angioma CNS HB Pancreatic NETs Low risk for PHEO Low risk for RCC 	<ul style="list-style-type: none"> PHEO Retinal angioma CNS HB Low risk for RCC 	<ul style="list-style-type: none"> PHEO Retinal angioma CNS HB Pancreatic cysts Pancreatic NETs RCC 	<ul style="list-style-type: none"> PHEO CNS HB Pancreatic NETs (rare)
Most common <i>VHL</i> variants	Truncating variants, exon deletions	Gene deletions encompassing <i>VHL</i>	Missense variants (e.g., p.Y98H, p.Y112H, p.V116F)	Missense variants (e.g., p.R167Q, p.R167W)	Missense variants (e.g., p.V84L, p.L188V)

NOTE: Based on ref. 6.

Abbreviation: CNS HB, central nervous system hemangioblastoma.

mutation type (Table 1). The clinical diagnosis is confirmed upon identification of a pathogenic variant in the *VHL* gene, the gold standard for vHL diagnosis (6).

Although many of the lesions associated with vHL present in the third and fourth decades of life, the age range of initial manifestations is wide (Table 2), and children are particularly vulnerable, as pediatric patients are at risk of developing hemangioblastomas and PHEO that can remain clinically occult until symptoms become severe. Moreover, when vHL-related tumors do occur, these malignancies can cause profound and lasting consequences (6–11).

Although most vHL-related tumors are histologically benign, morbidity due to mass effect can be significant (e.g., vision loss due to retinal hemangioblastoma; ref. 12). In addition, surgical intervention for tumors may be associated with operative/post-operative complications, such as hemorrhage from a large CNS hemangioblastoma (13). Historically, substantial mortality was attributable to RCC, pancreatic NET, and CNS hemangioblastoma (11). Fortunately, these risks have been mitigated in recent years by the institution of comprehensive surveillance paradigms, leading to early recognition of tumors and their multidisciplinary management (14–16).

Molecular genetics of vHL

vHL results from pathogenic variants in the *VHL* gene (6). vHL is inherited in an autosomal dominant pattern. Approximately 80% of individuals with vHL disease have an affected parent, and

about 20% result from a *de novo* pathogenic variant. Genetic testing is indicated in first-degree relatives of individuals with pathogenic variants in *VHL*, as well as any child diagnosed with any of the following:

- Retinal angioma (hemangioblastoma)
- CNS hemangioblastoma
- Clear cell RCC
- PHEO or paraganglioma
- ELST
- Epididymal or adnexal papillary cystadenoma
- Multiple pancreatic cysts or pancreatic NET
- Multiple renal cysts

The type of variant identified in the *VHL* gene has been shown to account for differences in PHEO risk, with a strong genotype-phenotype correlation (Table 1; ref. 17). Truncating variants or exon deletions in the *VHL* gene are identified among individuals with vHL type I and confer a relatively low risk of PHEO (17, 18). In contrast, vHL type II is associated with missense variants that generally do not affect the protein structure and are associated with a relatively higher risk of PHEO (17).

The risk for RCC and hemangioblastoma in affected individuals may reflect the ability of the variant protein to regulate the hypoxia-inducible factor (HIF) pathway (19, 20). Higher HIF expression appears to result in lower risk of RCC and hemangioblastoma. In addition, specific missense variants in *VHL* are also associated with hereditary polycythemia. Most of these variants lead solely to the polycythemia phenotype; however, a few variants (L188V, G104V, V130I, and Y175C) have been associated with both polycythemia and PHEO (21, 22).

Table 2. Lifetime risks of vHL-associated tumors

Tumor ^a	Risk	Youngest/mean age of diagnosis (years)
CNS hemangioblastoma	60%–80%	9/30
Cerebellar	44%–72%	9/31
Brainstem	10%–25%	12/32
Spinal	13%–50%	8/33
Retinal angioma/hemangioblastoma	25%–60%	0/25
Renal	25%–75%	12/39
Cyst	42%	12/37
RCC ^b	17%–70%	13/44
PHEO ^c	10%–25%	2/27
ELST ^d	10%–15%	6/22
Pancreatic	35%–75%	5/36
Cyst	21%	5/33
NET ^e	10%–17%	16/35
Papillary cystadenoma		
Epididymis	25%–60%	17/24
Broad ligament ^f	10%	16/unknown (16–46)

^aReferences 6, 23, and 24.

^bReferences 7, 11, 31, and 70.

^cReference 10.

^dReference 8.

^eReference 9.

^fReference 6.

Cancer screening/surveillance protocols

Early recognition and testing of at-risk individuals is key to the prevention of morbidity and mortality in vHL. Families with a history of vHL should be counseled regarding the importance of identifying at-risk children, performing genetic testing to identify mutation carriers, and initiating periodic surveillance. For example, children are susceptible to visual loss as result of retinal hemangiomas that, in the absence of surveillance, may go unnoticed until retinal damage becomes severe. CNS hemangioblastomas, if detected early, may be surgically excised with minimal damage to surrounding tissue.

Lifelong surveillance is, therefore, mandated given ongoing risks for tumor development and malignant growth with increasing age, and evidence that longitudinal surveillance in vHL limits morbidity and mortality (14–16). Multiple groups have developed surveillance guidelines (Table 3). Historically, these guidelines were based on expert consensus, with early screening targeted to tumor spectrum, risk levels, and typical ages of presentation (7, 23–25). Recent efforts have employed mathematical modeling methodologies, informed by the available clinical data,

Table 3. Existing paradigms for vHL tumor surveillance

Tumor	VHL Alliance 2015 ^a	Binderup et al. 2013 ^b (Denmark)	Hes et al. 2001 ^c (the Netherlands)	Kruizinga et al. 2014 ^d
Retinal HB	Annual eye exam ≥ 1 y	Annual eye exam ≥ 0 y	Annual eye exam ≥ 5 y	Biennial eye exam ≥ 7 y
PHEO	Annual PFM or UFM ≥ 5 y	Annual PFM ≥ 5 y	Annual PFM and UFM ≥ 10 y (serum and 24-h urine)	Every 4 y screen ≥ 0 y
ELST	Annual abd U/S 8–15 y Annual abd imaging ≥ 16 y (alternate U/S and MRI)	Q 1 y audiology eval ≥ 5 y	Q 1 y abd U/S 10–14 y Q 1 y abd imaging ≥ 15 y (alternate U/S and MRI)	None specified
CNS HB	Q 2–3 y audiology eval ≥ 15 y Biennial MRI b/s ≥ 16 y	MRI b/s once 8–14 y Biennial MRI b/s ≥ 15 y	Biennial MRI b/s ≥ 15 y	Annual MRI b/s ≥ 14 y
RCC	Annual abd U/S 8–15 y	Annual abd imaging ≥ 15 y (U/S or MRI)	Annual abd U/S 10–14 y	Annual screen ≥ 18 y
Pancreatic NET	Annual abd imaging ≥ 16 y (alternate U/S and MRI)	Annual abd imaging ≥ 15 y (U/S or MRI)	Annual abd imaging ≥ 15 y (alternate U/S and MRI)	Biennial screen ≥ 16 y
	Annual abd U/S 8–15 y		Annual abd U/S 10–14 y	
	Annual abd imaging ≥ 16 y (alternate U/S and MRI)		Annual abd imaging ≥ 15 y (alternate U/S and MRI)	

Abbreviations: abd, abdominal; eval, evaluation; h, hour; HB, hemangioblastoma; MRI b/s = MRI brain/spine; PFM, plasma-free metanephrines; Q, every; UFM, urinary fractionated metanephrines; U/S, ultrasound; y, year/years.

^aReference 23.

^bReference 24.

^cReference 7.

^dReference 26.

to predict the ideal ages to initiate surveillance and the most appropriate frequency for ongoing screening (26).

Our current consensus recommendations for screening of vHL-related tumors incorporate the elements of existing paradigms into a new surveillance regimen, with an emphasis on pediatric patients (Table 4). Broadly, the age-specific tumor risk, the youngest reported ages of occurrence, presumed growth rate, and the potential clinical impact of tumor progression were all considered in developing these new surveillance recommendations based on risk in children and adolescents.

Regardless of age, every individual with vHL should undergo an annual history and physical examination, including blood pressure assessment and a comprehensive neurologic evaluation, assessing for deficits including evidence of visual disturbance or hearing impairment. Ideally, these visits would be conducted by a medical provider experienced in caring for individuals with vHL, who has access to a multidisciplinary team with expertise in managing vHL-associated tumors. At these visits, education should also be provided on the signs and symptoms that could raise concern.

Beginning from birth, ophthalmology exams should be performed annually, with particular attention to the retina to monitor for retinal hemangioblastomas. This is consistent with most current screening paradigms. These exams should be conducted by an ophthalmologist experienced in pediatric retinal evaluation. In addition, starting at 2 years of age, PHEO surveillance should commence with blood pressure checks at every medical visit (using standard tables based on age and height; ref. 27), and with annual plasma or urine metanephrine levels. This screening interval is in agreement with standard practices, but the youngest age to initiate PHEO screening advocated in existing screening protocols is 5 years of age. There are multiple reports of PHEO occurring in children at younger ages, driving the impetus to advance the onset of screening (10, 26). We note that younger patients with PHEO bore missense variants (type II/III) rather than truncating or deletion variants (type I). This may inform future iterations of recommendations with respect to the age to

initiate PHEO surveillance. In the course of surveillance, attention should be given to avoiding potentially interfering foods and medications that may confound interpretation of biochemical testing (Supplementary Table S1).

By 5 years of age, biennial audiology evaluations should be commenced to screen for ELST. This is in agreement with current screening paradigms. By 8 years of age, biennial MRI of the brain and spine should begin to monitor for CNS hemangioblastomas. Our recommendation to start this longitudinal surveillance at 8 years is significantly younger than dictated by other guidelines. Although the risk of CNS hemangioblastomas prior to adolescence is relatively low, multiple instances have been reported earlier in childhood (6, 23, 24). Moreover, these tumors may cause substantial morbidity with progression, particularly those lesions associated with peritumoral cysts that are more commonly found in younger individuals with vHL and that may progress more rapidly (28). As the risk of CNS hemangioblastomas rises in adolescence, and given the approximately 7% risk of hemangioblastomas developing in the interval between every-other-year MRIs, consideration may be given to increasing the screening frequency to annually starting in mid-adolescence, in contrast to the biennial surveillance recommended in established guidelines (4). The risk of rapidly growing hemangioblastomas, however, needs to be better quantified before this approach can be formally recommended. Cranial imaging should include thin cuts through the internal auditory canals, as a complement to audiology evaluations for ELST screening due to the possibility of ELST detection prior to development of audio-vestibular symptoms (29). Because spinal hemangioblastomas can occur at any segment of the spinal cord, it should be emphasized that spinal imaging should extend the entire length of the cord and not be restricted to only the cervical cord (30).

Our proposed surveillance for visceral manifestations of vHL diverges significantly from established recommendations. Primary screening for PHEO with imaging is not advocated given the high sensitivity of biochemical screening measures. Therefore, the characteristics of RCC (31) and pancreatic NET (6, 9, 23, 24)

Table 4. Proposed vHL tumor surveillance regimen with an emphasis on the pediatric age range

Tumor	Recommended surveillance	Age to begin	Interval
Retinal HB	Eye exam including retina ^a	Birth	Annual
PHEO	Blood pressure at all medical visits ^b	2 years	
	PFM ^{c,d,e,f} or 24-h urine fractionated metanephrines ^g	2 years	Annual
ELST	Audiogram	5 years	Biennial
CNS HB	MRI brain with and without contrast ^h	8 years	Biennial ⁱ
	MRI spine with contrast		
RCC	MRI abdomen ^j	10 years	Annual
Pancreatic NET	MRI abdomen	10 years	Annual

Abbreviations: HB, hemangioblastoma; PFM, plasma-free metanephrines.

^aDuring childhood, ophthalmologic examination should be performed by an ophthalmologist with experience in pediatric retinal examination.

^bBlood pressure in children should be assessed using age- and height-specific normative ranges (https://www.nhlbi.nih.gov/files/docs/guidelines/child_tbl.pdf).

^cReference to pediatric reference intervals for plasma (71, 72) and urine (73) metanephrines should be considered.

^dIdeally, to limit false positive results, PFMs should be collected from an indwelling venous catheter after patient has been lying supine for ≥ 30 minutes. Clinicians may elect to bypass this approach, but marginally elevated results should prompt repetition of testing under ideal conditions.

^eSeveral foods and medications may interfere with metanephrine analysis and should be avoided prior to testing. These are summarized in Supplementary Table S1.

^fRecommended action based on plasma metanephrines:

– Confirm interfering agents were avoided prior to testing (Supplementary Table S1).

– If $\geq 4\times$ upper limit of reference range: consistent with disease, proceed with imaging to localize lesion.

– If $2\times$ – $4\times$ upper limit of reference range: repeat testing in 2 months.

– If marginally elevated: repeat testing in 6 months or consider clonidine suppression test to exclude false positivity (74, 75).

^gTwenty-four-hour urine fractionated metanephrines are an acceptable alternative to plasma metanephrines once patients are continent of urine.

^hIncluding thin cuts through internal auditory canals.

ⁱSome providers may choose to advance the frequency of CNS imaging to annually starting in adolescence due to retrospective data describing relatively quick growing hemangioblastomas in between every-2-year scans.

^jMRI surveillance for RCC and pancreatic NET may be part of the same study, provided that dedicated renal sequences are included.

should drive the timing of screening initiation and the interval of ongoing surveillance. We recommend that surveillance for visceral manifestations of vHL should be implemented at 10 years of age, with annual MRI of the abdomen to maximize the sensitivity and maintain the consistency of the screening methodology for RCC and pancreatic NET detection. To ensure optimal detection for RCC, this abdominal MRI should be performed per institutional protocol used for renal evaluation. Ultrasound may be considered to complement the MRI, whereas CT is reserved for rare circumstances where biochemical abnormalities are detected and MRI is contraindicated. The latter places high value on reduction of exposure to ionizing radiation, in accordance with the position of the VHL Alliance (23).

Although longstanding evidence supports genotype–phenotype correlations in vHL, these are generally not considered in tailoring screening recommendations, as the correlations are still being characterized. This may change as data accrued from systematic surveillance become available. It should be recognized that the tumor surveillance in vHL is time-consuming and may incur substantial financial and psychosocial burdens (32), as discussed further in the genetic counseling article in

this CCR Pediatric Oncology Series (33). However, these burdens may be diminished by experienced multidisciplinary teams through care coordination and enhanced education.

Conclusions

We propose a tumor surveillance paradigm for individuals with vHL based on specific risks in childhood and adolescence. Although these recommendations derive from existing paradigms, in considering screening onset and intervals, we placed high priority on the earliest ages of tumor onset, potential tumor growth rates, and the clinical impact of delayed detection of these tumors in children and adolescents. We have not made specific recommendations for adults with vHL, and practitioners should rely on the existing screening regimens for adults with vHL (Table 3).

Despite our reliance on available data, these guidelines remain largely based on expert opinion. The next logical step would incorporate a prospective assessment of clinical outcomes of individuals with vHL, screened according to these proposed guidelines. In addition, future advances in early detection methodologies may be realized through identification of reliable biomarkers for the aberrant vascular proliferation occurring in vHL. Finally, no measures are currently available to prevent individuals from developing vHL manifestations. Future investigations, however, could include the identification and application of strategies to inhibit aberrant vascular growth (34). The main impediments to all of these avenues of future study are the relatively low prevalence of vHL and the prolonged duration over which associated tumors may arise. These characteristics of the condition make accrual of sufficient numbers of affected patients, and/or samples, for these studies challenging and will depend on collaborative multi-institutional efforts.

Hereditary PHEO/Paraganglioma Syndromes

Introduction

Hereditary paraganglioma and PHEO syndromes (HPP) are characterized by rare and usually benign tumors of neural crest origin that are symmetrically distributed along the paravertebral axis from the base of the skull and neck to the pelvis. In addition to paraganglioma/PIIEOs, patients with IIPP syndromes can develop renal cancers, gastrointestinal stromal tumors (GIST), pituitary adenomas, and other rare tumor types. The genes in which pathogenic variants are known to cause HPP syndromes collectively include the *SDHx* genes, a group of multiple nuclear genes encoding subunits of the succinate dehydrogenase (SDH) enzyme complex. This enzyme complex catalyzes the conversion of succinate to fumarate in the Krebs cycle and serves as complex II of the electron transport chain. *SDHx* genes include SDH subunits A to D and SDH assembly factor 2 (SDHAF2), which is a stabilizing protein required for the flavination of SDHA (35). Other non-*SDHx* genes associated with hereditary paraganglioma/PHEOs include the *MAX* gene encoding a member of the helix-loop-helix leucine zipper family of transcription factors that regulates cell proliferation, *TMEM127* that encodes a transmembrane protein that negatively regulates mTOR (36, 37). Recently HIF2 α , EGLN1, and KIF1 β have been implicated in the development of PHEO/paraganglioma (38, 39). It should also be borne in mind that PHEO/paraganglioma are components of other hereditary tumor predisposition syndromes associated with the *RET*, *VHL*, *NFI*, and *FH* genes, described elsewhere.

As with vHL, although the tumors associated with HPP syndrome are most frequently histologically benign, they can result in significant clinical morbidity related to mass effect, cranial nerve palsies, or hypertension/tachycardia resulting from catecholamine excess. If left untreated, a subset of these tumors will metastasize. A high malignant potential has been specifically recognized for *SDHB*-related tumors and is associated with tumor size at the time of diagnosis (40, 41). The increased metastatic potential and aggressive nature of *SDHx*-related paragangliomas compared with *de novo* paragangliomas without underlying germline predisposition must be taken into account when developing early tumor surveillance in patients with HPP.

Paragangliomas in the skull base, neck, and upper mediastinum are primarily associated with the parasympathetic nervous system and generally do not hypersecrete catecholamine or other hormones, although a subset can secrete dopamine. In contrast, paragangliomas in the lower mediastinum, abdomen, and pelvis are typically associated with the sympathetic nervous system and usually hypersecrete catecholamines. PIIEOs are catecholamine-secreting paragangliomas confined to the adrenal medulla.

The most current studies suggest that up to 35% of PHEO/paraganglioma are hereditary (42). The diagnosis of HPP syndrome is based on molecular genetic testing, which should be offered to all patients with paraganglioma or PHEO. However, recognition of bilateral paraganglioma/PHEO and/or a strong autosomal dominant familial presentation of paraganglioma/PHEO may also lead to a clinical diagnosis of HPP. Assessment of the family history should include specific inquiry regarding relatives with sudden death, and it should detail the spectrum of tumors associated with HPP syndrome, some of which are known by different names (paragangliomas, e.g., may be called glomus tumors or extra-adrenal PHEOs). The medical history should include symptoms of catecholamine excess, such as elevation in blood pressure, headaches, diaphoresis, and palpitations, and also symptoms attributable to mass effect, such as dysphagia, hearing loss, and dysarthria. The exam should include assessment for arrhythmias and palpable masses (35).

GISTs occurring in the setting of pathogenic *SDHx* germline variants have distinct clinical features. The *SDHx*-related GISTs are almost always located in the stomach, arising from the interstitial cells of Cajal in a submucosal location. Multifocal gastric tumors are common. Because of their gastric location, by far the most common presentation for *SDHx*-related GISTs is gastric bleeding.

Molecular genetics of HPP syndromes

Pathogenic missense and truncating variants in *SDH B, C*, and *D* underlie the majority of patients with HPP syndromes (after excluding other syndromes such as vHL, multiple endocrine neoplasia and, less frequently, neurofibromatosis and hereditary leiomyomatosis and RCC). Variants in *SDHA* and *SDHAF2*, albeit rare, have also been reported. Although *SDHx*-related GISTs can be due to many of the different *SDH* genes (including *SDHC* epimutations), the *SDHA* germline mutations seem to be the most common germline mutations associated with GISTs (43). In the setting of tumor susceptibility, the *SDHx* genes act as tumor suppressors. The tumorigenesis mediated by mutated *SHDx* genes is thought to result from elevations in cellular succinate concentrations that stabilize the HIF α transcription factor by inhibiting prolyl hydroxylases, thus preventing HIF α degradation by ubiquitination (44). High levels of HIF α are strongly implicated in

promoting tumor growth and metastasis through the role of HIF α in initiating angiogenesis and regulating cellular metabolism to overcome hypoxia (45). In addition, several reports suggest that *SDHx*-related tumors display a hypermethylator phenotype associated with downregulation of key genes involved in neuroendocrine differentiation (46, 47). *TMEM127* and *MAX* also function as tumor suppressors, and loss of heterozygosity has been demonstrated in PHEO/paraganglioma tumor tissue.

Susceptibility to paraganglioma/PHEO, as well as other associated tumors, is either inherited in a strict autosomal dominant fashion (*SDHA*, *SDHB*, *SDHC*, and *TMEM127*) or in an autosomal dominant fashion modified by parent of origin. *SDHD*, *SDHAF2*, and *MAX* variants present with this parent-of-origin-dependent tumorigenesis, wherein tumor formation occurs almost exclusively in the context of paternal transmission of the variant due to maternal imprinting (i.e., affected children of carrier fathers will present with disease, but affected children of carrier mothers will not).

Penetrance estimates have been defined for variants in *SDHD* and *SDHB*. For *SDHD* variants, the penetrance is about 90% for probands and relatives identified through variant testing (48). Penetrance estimates for *SDHB* variant carriers have been more variable due to ascertainment and analytic differences between studies. Early studies estimated an approximately 77% risk for paraganglioma by age 70 (49); however, more recent estimates based on testing of extended families have estimated the lifetime risk to be between 30% and 50% (50) and may be even as low as 14% by age 60 (51). Variants in *SDHA*, *SDHAF2*, and *SDHC* are rare, and specific penetrance estimates have not been calculated. A single-center review of eight probands identified with *SDHC* variants noted that none of them had a family history of paraganglioma, suggesting that the penetrance is likely incomplete (52). *SDHAF2* variants have been described in a large Dutch family. In this kindred, 11 of 16 individuals (69%) who had paternally inherited the variant were found to have paraganglioma (50, 53). Cumulative penetrance of *TMEM127*-associated PHEOs has been estimated to be 32% from a large family with 34 members (54).

Patients with *SDHB* variants have a positive family history in 33% of cases, present with single tumors at a mean age of 25 to 30 years (range, 6–77), and are strongly associated with extra-adrenal sympathetic paragangliomas, mainly in the abdomen and pelvis. About 20% may also have PHEOs, and, as discussed, paragangliomas in these patients have a substantial propensity to metastasize (25, 55–59). Paraganglioma development in *SDHB* variant carriers is generally associated with higher morbidity and mortality than pathogenic variants in the other *SDHx* genes due to the greater risk for malignancy and metastasis. In comparison with individuals with sporadic, malignant paraganglioma, those with malignant disease and a germline *SDHB* variant have shorter survival (55). Although less common than malignant extra-adrenal sympathetic paragangliomas, malignant PHEOs do occur and may be more common among individuals with a germline variant in *SDHB* compared with other PHEO-predisposing loci.

Patients with *SDHD*, *SDHC*, and *SDHAF2* variants are more frequently associated with parasympathetic skull-base and neck paragangliomas (49, 52, 53). Individuals with *SDHD* variants present at a mean age of 28 to 31 years (range, 12–70) with multiple tumors, whereas those with *SDHC* variants most commonly develop a solitary tumor (about 77%), with a mean age of

Table 5. Proposed HPP surveillance regimen

Tumor	Recommended surveillance	Age to begin	Interval
PGL/PHEO	Blood pressure at all medical visits ^a	6–8 years	Annual (at minimum)
	Plasma methoxytyramine	6–8 years	Annual
	PFM ^{b,c,d,e} or 24-h urine fractionated metanephrines ^f	6–8 years	Annual
	Optional: serum chromogranin A	6–8 years	Annual
	Whole-body MRI (skull base to pelvis) ^g	6–8 years	Biennial
	Optional: neck MRI ± contrast ^h	6–8 years	Biennial
GIST	Complete blood count (w/RBC indices)	6–8 years	Annual

Abbreviations: PFM, plasma-free metanephrines; RBC, red blood cell.

^aBlood pressure in children should be assessed using age- and height-specific normative ranges (https://www.nhlbi.nih.gov/files/docs/guidelines/child_tbl.pdf, ref. 27).

^bReference to pediatric reference intervals for plasma (71, 72) and urine (73) metanephrines should be considered.

^cIdeally, to limit false positive results, PFM should be collected from an indwelling venous catheter after patient has been lying supine for ≥ 30 minutes. Clinicians may elect to bypass this approach, but marginally elevated results should prompt repetition of testing under ideal conditions.

^dSeveral foods and medications may interfere with metanephrine analysis and should be avoided prior to testing. These are summarized in Supplementary Table S1.

^eRecommended action based on plasma metanephrines:

– Confirm interfering agents were avoided prior to testing (Supplementary Table S1).

– If $\geq 4\times$ upper limit of reference range: consistent with disease, proceed with imaging to localize lesion.

– If $2\times$ – $4\times$ upper limit of reference range: repeat testing in 2 months.

– If marginally elevated: repeat testing in 6 months or consider clonidine suppression test to exclude false positivity (74, 75).

^fTwenty-four-hour urine fractionated metanephrines are an acceptable alternative to plasma metanephrines once patients are continent of urine.

^gSpecific attention also paid to the kidneys due to rare risk of RCC.

^hDepending on preferences of local radiologists, dedicated MRI of the neck may be preferred to inclusion in whole-body MRI. If this is the case, it should be performed concomitant with WBMRI.

diagnosis of 38 years of age (range, 15–40; refs. 25, 57, 58). Eighty percent of reported *SDHC*-related paragangliomas originate in the head and neck, and malignant disease has been rarely reported (2%; ref. 52). Only head and neck tumors, predominantly carotid body tumors (70%), have been reported in *SDHAF2* variant carriers, with an average age of onset of 33 years (range, 22–47). However, all descriptions of the *SDHAF2* phenotype are based on a single family (53, 60). Germline *SDHA* variants have been observed in association with both PHEOs and paragangliomas (sympathetic and parasympathetic; ref. 60).

Screening/Surveillance

Tumorigenesis is rare in the first decade of life among individuals predisposed to HPP. The youngest patient that we have identified with paraganglioma was 6 years at diagnosis (59). After reviewing the literature and clinical practice across many different institutions and countries, we recommend initiating tumor surveillance at 6 to 8 years of age. Although we acknowledge specific genetic lesions drive varying clinical presentations, we do not think that the genotype–phenotype relationships are currently well enough defined to justify distinct surveillance paradigms based on genotype. Therefore, we advocate a single regimen for all carriers of variants associated with HPP syndromes, with a need to accumulate enough data over time to make more gene-specific recommendations.

Surveillance recommendations for IIPP syndromes resemble those for vHL with respect to PHEO/paraganglioma screening (Table 5). However, we recognize the existence of silent (non-secretory) paragangliomas, particularly those occurring in the head and neck, which occur with greater frequency than in vHL and, thus, emphasize the importance of radiologic surveillance in HPP syndrome in addition to biochemical screening. This screening will also facilitate identification of RCCs, which have rarely been associated with germline *SDHx* variants. Although limited data exist, whole-body imaging was recently studied among individuals with *SDH*-associated paraganglioma syndromes and identified six tumors among asymptomatic carriers. Sensitivity and specificity of whole-body imaging were

87.5% and 94.7%, respectively, whereas biochemical surveillance yielded sensitivity and specificity of 37.5% and 94.5%, respectively (61). Although there is increasing interest in the use of contrast-enhanced ultrasound (CEUS; ref. 62), there are insufficient data to recommend CEUS over MRI for routine surveillance. As some paragangliomas are predominantly dopaminergic (particularly those associated with *SDHB*, *C*, *D*), plasma methoxytyramine analysis is advocated in addition to metanephrines (63–65). Finally, chromogranin A, a NET marker, has been demonstrated to enhance detection of PHEO/paragangliomas by 22%, with minimal impact on specificity, and can be included in annual surveillance (66).

There are currently no data to support routine imaging for GISTs in at-risk patients. Formal studies of the sensitivity of whole-body MRI for submucosal gastric masses have not been performed, but one would anticipate that the sensitivity will be relatively low for small GISTs. Imaging approaches more likely to detect submucosal gastric GISTs include CT with oral and intravenous contrast, with or without FDG-PET. Both of these imaging modalities carry the risk of exposure to radiation. Given the relatively low incidence of GIST in HPP, the benefits are unlikely to outweigh the risks of exposure to radiation and follow-up of false positives. On the basis of this information, we do not advocate imaging surveillance specifically for GIST detection in patients with HPP syndrome. Because essentially all GISTs in IIPP patients are gastric (in some series, 95% of patients present with gastric bleeding and/or anemia, and in some cases, GISTs can present with rapid bleeding and significant anemia; refs. 67–69), based solely on a consensus of opinion, we recommend an annual complete blood count to screen for incident anemia during surveillance. The effectiveness of this approach and the frequency of incidental findings requiring further evaluation with hematology referral or endoscopy are not known. There are insufficient data to define the typical age of onset of GIST in this population, so we propose that the age of onset and the interval for surveillance is the same as for PHEO/paraganglioma (at least until such time as future evidence dictates otherwise).

Conclusions

The identification of the genetic basis of the HPP syndromes has led to significant advances in clinical care for these patients, providing prognostic insights as well as opportunities for early detection and treatment of component tumors. Further research is needed to clarify how individual genotypes can more reliably predict phenotypic behavior, and whether and how surveillance should differ based on genotype. In addition, there remains an urgent and unmet clinical need to develop improved therapies for patients with metastatic PHEO/paragangliomas.

Disclosure of Potential Conflicts of Interest

W. K. Kohlmann reports receiving commercial research grants from Myriad. No potential conflicts of interest were disclosed by the other authors.

Grant Support

G.E. Tomlinson was supported by the NCI under award number 5P30CA054174-21.

Received February 27, 2017; revised April 24, 2017; accepted April 27, 2017; published online June 15, 2017.

References

- Chittiboia P, Lonser RR. Von Hippel-Lindau disease. *Handb Clin Neurol* 2015;132:139–56.
- Hippel Ev. Ueber eine sehr seltene Erkrankung der Netzhaut. *Albrecht von Graefes Arch Ophthalmol* 1904;59:83–106.
- Lindau A. Zur Frage der Angiomatosis Retinae und Ihrer Hirncomplication. *Acta Ophthalmol* 1927;4:193–226.
- Poulsen ML, Budtz-Jorgensen E, Bisgaard ML. Surveillance in von Hippel-Lindau disease (vHL). *Clin Genet* 2010;77:49–59.
- Lonser RR, Glenn GM, Walther M, Chew EY, Libutti SK, Linehan WM, et al. von Hippel-Lindau disease. *Lancet* 2003;361:2059–67.
- Nielsen SM, Rhodes L, Blanco I, Chung WK, Eng C, Maher ER, et al. Von Hippel-Lindau disease: genetics and role of genetic counseling in a multiple neoplasia syndrome. *J Clin Oncol* 2016;34:2172–81.
- Hes FJ, van der Luijt RB, Lips CJ. Clinical management of Von Hippel-Lindau (VHL) disease. *Neth J Med* 2001;59:225–34.
- Bausch B, Wellner U, Peyre M, Boedeker CC, Hies FJ, Anglani M, et al. Characterization of endolymphatic sac tumors and von Hippel-Lindau disease in the International Endolymphatic Sac Tumor Registry. *Head Neck* 2016;38Suppl 1:E673–9.
- Libutti SK, Choyke PL, Alexander HR, Glenn G, Bartlett DL, Zbar B, et al. Clinical and genetic analysis of patients with pancreatic neuroendocrine tumors associated with von Hippel-Lindau disease. *Surgery* 2000;128:1022–7.
- Sovinz P, Urban C, Uhrig S, Stepan V, Lackner H, Schwinger W, et al. Pheochromocytoma in a 2.75-year-old-girl with a germline von Hippel-Lindau mutation Q164R. *Am J Med Genet A* 2010;152A:1752–5.
- Maher ER, Yates JR, Harries R, Benjamin C, Harris R, Moore AT, et al. Clinical features and natural history of von Hippel-Lindau disease. *Q J Med* 1990;77:1151–63.
- Feletti A, Anglani M, Scarpa B, Schiavi F, Boaretto F, Zovato S, et al. Von Hippel-Lindau disease: an evaluation of natural history and functional disability. *Neuro Oncol* 2016;18:1011–20.
- Wind JJ, Lonser RR. Management of von Hippel-Lindau disease-associated CNS lesions. *Expert Rev Neurother* 2011;11:1433–41.
- Binderup ML, Jensen AM, Budtz-Jorgensen E, Bisgaard ML. Survival and causes of death in patients with von Hippel-Lindau disease. *J Med Genet* 2017;54:11–8.
- Priesemann M, Davies KM, Perry LA, Drake WM, Chew SL, Monson JP, et al. Benefits of screening in von Hippel-Lindau disease—comparison of morbidity associated with initial tumours in affected parents and children. *Horm Res* 2006;66:1–5.
- Prasad R, Johnston LB, Savage MO, Martin L, Perry LA, Storr HL. Pediatric endocrine screening for von Hippel-Lindau disease: benefits and the challenge of compliance. *J Endocrinol Invest* 2011;34:296–9.
- Nordstrom-O'Brien M, van der Luijt RB, van Rooijen E, van den Ouweland AM, Majoor-Krakauer DF, Lolkema MP, et al. Genetic analysis of von Hippel-Lindau disease. *Hum Mutat* 2010;31:521–37.
- Stebbins CE, Kaelin WG Jr, Pavletich NP. Structure of the VHL-ElonginC-ElonginB complex: implications for VHL tumor suppressor function. *Science* 1999;284:455–61.
- Binderup ML, Budtz-Jorgensen E, Bisgaard ML. Risk of new tumors in von Hippel-Lindau patients depends on age and genotype. *Genet Med* 2016;18:89–97.
- Gossage L, Eisen T, Maher ER. VHL, the story of a tumour suppressor gene. *Nat Rev Cancer* 2015;15:55–64.
- Gossage L, Pires DE, Olivera-Nappa A, Asenjo J, Bycroft M, Blundell TL, et al. An integrated computational approach can classify VHL missense mutations according to risk of clear cell renal carcinoma. *Hum Mol Genet* 2014;23:5976–88.
- Capodimonti S, Teofili L, Martini M, Cenci T, Iachinoto MC, Nuzzolo ER, et al. Von hippel-lindau disease and erythrocytosis. *J Clin Oncol* 2012;30:e137–9.
- VHL Alliance. The VHL handbook. 5th ed. Boston (MA): VHL Alliance; 2015.
- Binderup ML, Bisgaard ML, Harbud V, Moller HU, Gimsing S, Friis-Hansen L, et al. Von Hippel-Lindau disease (vHL). National clinical guideline for diagnosis and surveillance in Denmark. 3rd edition. *Dan Med J* 2013;60:B4763.
- Lefebvre M, Foulkes WD. Pheochromocytoma and paraganglioma syndromes: genetics and management update. *Curr Oncol* 2014;21:e8–17.
- Kruizinga RC, Sluiter WJ, de Vries EC, Zonnenberg BA, Lips CJ, van der Horst-Schrivers AN, et al. Calculating optimal surveillance for detection of von Hippel-Lindau-related manifestations. *Endocr Relat Cancer* 2014;21:63–71.
- National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004;114:555–76.
- Huntoon K, Wu T, Elder JB, Butman JA, Chew EY, Linehan WM, et al. Biological and clinical impact of hemangioblastoma-associated peritumoral cysts in von Hippel-Lindau disease. *J Neurosurg* 2016;124:971–6.
- Poulsen ML, Gimsing S, Kosteljanetz M, Moller HU, Brandt CA, Thomsen C, et al. von Hippel-Lindau disease: surveillance strategy for endolymphatic sac tumors. *Genet Med* 2011;13:1032–41.
- Lonser RR, Butman JA, Huntoon K, Asthagiri AR, Wu T, Bakhtian KD, et al. Prospective natural history study of central nervous system hemangioblastomas in von Hippel-Lindau disease. *J Neurosurg* 2014;120:1055–62.
- Ferragut J, Caimari M, Rituerto B, Gomez-Rivas B, Herrera M, Alonso F. Pheochromocytoma and clear-cell renal carcinoma in a child with von Hippel-Lindau disease: a patient report. *J Pediatr Endocrinol Metab* 1999;12:579–82.
- Lammens CR, Aaronson NK, Hes FJ, Links JP, Zonnenberg BA, Lenders JW, et al. Compliance with periodic surveillance for Von-Hippel-Lindau disease. *Genet Med* 2011;13:519–27.
- Druker H, Zellek K, McGee RB, Scollon S, Kohlmann W, Schneider KA, et al. Genetic counselor recommendations for cancer predisposition evaluation and surveillance in the pediatric oncology patient. *Clin Cancer Res* 2017;23:doi: 10.1158/1078-0432.CCR-17-0834.
- Migliorini D, Haller S, Merker D, Pugliesi-Rinaldi A, Koka A, Schaller K, et al. Recurrent multiple CNS hemangioblastomas with VHL disease treated with pazopanib: a case report and literature review. *CNS Oncol* 2015;4:387–92.
- Kirmani S, Young WF. Hereditary paraganglioma-pheochromocytoma syndromes. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LH, et al., editors. *GeneReviews* [Internet]. Seattle (WA): University of Washington, Seattle; 1993. 2008 May 21 [updated 2014 Nov 6].

36. Comino-Mendez I, Gracia-Aznarez FJ, Schiavi F, Landa I, Leandro-Garcia LJ, Leton R, et al. Exome sequencing identifies MAX mutations as a cause of hereditary pheochromocytoma. *Nat Genet* 2011;43:663–7.
37. Qin Y, Yao L, King EE, Buddavarapu K, Lenci RE, Chocron ES, et al. Germline mutations in TMEM127 confer susceptibility to pheochromocytoma. *Nat Genet* 2010;42:229–33.
38. Zhuang Z, Yang C, Lorenzo F, Merino M, Fojo T, Kebebew E, et al. Somatic HIF2A gain-of-function mutations in paraganglioma with polycythemia. *N Engl J Med* 2012;367:922–30.
39. Schlisio S, Kenchappa RS, Vredeveld LC, George RE, Stewart R, Greulich H, et al. The kinesin KIF1Bbeta acts downstream from EglN3 to induce apoptosis and is a potential 1p36 tumor suppressor. *Genes Dev* 2008;22:884–93.
40. Assadipour Y, Sadowski SM, Alimchandani M, Quezado M, Steinberg SM, Nilubol N, et al. SDHB mutation status and tumor size but not tumor grade are important predictors of clinical outcome in pheochromocytoma and abdominal paraganglioma. *Surgery* 2017;161:230–9.
41. Turkova H, Prodanov I, Maly M, Martucci V, Adams K, Widimsky J Jr, et al. Characteristics and outcomes of metastatic SDHB and Sporadic Pheochromocytoma/Paraganglioma: an National Institutes of Health Study. *Endocr Pract* 2016;22:302–14.
42. Martins R, Bugalho MJ. Paragangliomas/Pheochromocytomas: clinically oriented genetic testing. *Int J Endocrinol* 2014;2014:794187.
43. Boikos SA, Pappo AS, Killian JK, LaQuaglia MP, Weldon CB, George S, et al. Molecular subtypes of KIT/PDGFRA wild-type gastrointestinal stromal tumors: a report from the National Institutes of Health Gastrointestinal Stromal Tumor Clinic. *JAMA Oncol* 2016;2:922–8.
44. Cottlieb E, Tomlinson IP. Mitochondrial tumour suppressors: a genetic and biochemical update. *Nat Rev Cancer* 2005;5:857–66.
45. Mubjeesh NJ, Amir S. Hypoxia-inducible factor (HIF) in human tumorigenesis. *Histol Histopathol* 2007;22:559–72.
46. Letouze E, Martinelli C, Loriot C, Burnichon N, Abermil N, Ottolenghi C, et al. SDH mutations establish a hypermethylator phenotype in paraganglioma. *Cancer Cell* 2013;23:739–52.
47. Fishbein L, Leshchiner I, Walter V, Danilova L, Robertson AG, Johnson AR, et al. Comprehensive molecular characterization of pheochromocytoma and paraganglioma. *Cancer Cell* 2017;31:181–93.
48. Ricketts CJ, Forman JR, Rattenberry E, Bradshaw N, Lalloo F, Izatt L, et al. Tumor risks and genotype-phenotype-proteotype analysis in 358 patients with germline mutations in SDHB and SDHD. *Hum Mutat* 2010;31:41–51.
49. Neumann HP, Pawlu C, Peczkowska M, Bausch B, McWhinney SR, Muresan M, et al. Distinct clinical features of paraganglioma syndromes associated with SDHB and SDHD gene mutations. *JAMA* 2004;292:943–51.
50. Jaffi M, Whitworth J, Rattenberry E, Vialard L, Kilby G, Kumar AV, et al. Evaluation of SDHB, SDHD and VHL gene susceptibility testing in the assessment of individuals with non-syndromic pheochromocytoma, paraganglioma and head and neck paraganglioma. *Clin Endocrinol (Oxf)* 2013;78:898–906.
51. Eijkelenkamp K, Osinga TE, de Jong MM, Sluiter WJ, Dullaart RP, Links TP, et al. Calculating the optimal surveillance for head and neck paraganglioma in SDHB-mutation carriers. *Fam Cancer* 2017;16:123–30.
52. Else T, Marvin ML, Everett JN, Cruber SB, Arts HA, Stoffel EM, et al. The clinical phenotype of SDH1C-associated hereditary paraganglioma syndrome (PGL3). *J Clin Endocrinol Metab* 2014;99:E1482–6.
53. Kunst HP, Rutten MH, de Monnik JP, Hoefsloot LH, Timmers HJ, Marres HA, et al. SDH1AF2 (PGL2-SDH15) and hereditary head and neck paraganglioma. *Clin Cancer Res* 2011;17:247–54.
54. Toledo SP, Lourenco DM Jr, Sekiya T, Lucon AM, Baena ME, Castro CC, et al. Penetrance and clinical features of pheochromocytoma in a six-generation family carrying a germline TMEM127 mutation. *J Clin Endocrinol Metab* 2015;100:E308–18.
55. Amar L, Baudin E, Burnichon N, Peyrard S, Silveira S, Bertherat J, et al. Succinate dehydrogenase B gene mutations predict survival in patients with malignant pheochromocytomas or paragangliomas. *J Clin Endocrinol Metab* 2007;92:3822–8.
56. Van Nederveen FH, Dinjens WN, Korpershoek E, De Krijger RR. The occurrence of SDHB gene mutations in pheochromocytoma. *Ann N Y Acad Sci* 2006;1073:177–82.
57. Eisenhofer G, Timmers HJ, Lenders JWM, Bornstein SR, Tiebel O, Mannelli M, et al. Age at diagnosis of pheochromocytoma differs according to catecholamine phenotype and tumor location. *J Clin Endocrinol Metab* 2011;96:375–84.
58. Benn DE, Gimenez-Roqueplo A-P, Reilly JR, Bertherat J, Burgess J, Byth K, et al. Clinical presentation and penetrance of pheochromocytoma/paraganglioma syndromes. *J Clin Endocrinol Metab* 2006;91:827–36.
59. Burnichon N, Rohmer V, Amar L, Herman P, Lebouilleux S, Darrouzet V, et al. The succinate dehydrogenase genetic testing in a large prospective series of patients with paragangliomas. *J Clin Endocrinol Metab* 2009;94:2817–27.
60. Bausch B, Schiavi F, Ni Y, Welander J, Patocs A, Ngeow J, et al. Clinical characterization of the pheochromocytoma and paraganglioma susceptibility genes sdha, tmem127, max, and sdhaf2 for gene-informed prevention. *JAMA Oncol* 2017 Apr 6. [Epub ahead of print].
61. Jaspersen KW, Kohlmann W, Cammon A, Slack H, Buchmann L, Hunt J, et al. Role of rapid sequence whole-body MRI screening in SDH1-associated hereditary paraganglioma families. *Fam Cancer* 2014;13:257–65.
62. Meister M, Choyke P, Anderson C, Patel U. Radiological evaluation, management, and surveillance of renal masses in Von Hippel-Lindau disease. *Clin Radiol* 2009;64:589–600.
63. van Duinen N, Corssmit EP, de Jong WH, Brookman D, Kema IP, Romijn JA. Plasma levels of free metanephrines and 3-methoxytyramine indicate a higher number of biochemically active HNPGL than 24-h urinary excretion rates of catecholamines and metabolites. *Eur J Endocrinol* 2013;169:377–82.
64. Peitzsch M, Prejbisz A, Kroiss M, Beuschlein F, Arlt W, Januszewicz A, et al. Analysis of plasma 3-methoxytyramine, normetanephrine and metanephrine by ultra-performance liquid chromatography-tandem mass spectrometry: utility for diagnosis of dopamine-producing metastatic pheochromocytoma. *Ann Clin Biochem* 2013;50(Pt 2):147–55.
65. Gupta G, Pacak K; AACE Adrenal Scientific Committee. Precision medicine: an update on genotype-biochemical phenotype relationships in pheochromocytoma/paraganglioma patients. *Endocr Pract* 2017 Mar 23. [Epub ahead of print].
66. Zuber S, Wesley R, Prodanov T, Eisenhofer G, Pacak K, Kantorovich V. Clinical utility of chromogranin A in SDHx-related paragangliomas. *Eur J Clin Invest* 2014;44:365–71.
67. Zhang L, Smyrk TC, Young WF Jr, Stratakis CA, Carney JA. Gastric stromal tumors in Carney triad are different clinically, pathologically, and behaviorally from sporadic gastric gastrointestinal stromal tumors: findings in 104 cases. *Am J Surg Pathol* 2010;34:53–64.
68. De Vogelaere K, Aerts M, Haentjens P, De Greve J, Delvaux G. Gastrointestinal stromal tumor of the stomach: progresses in diagnosis and treatment. *Acta Gastroenterol Belg* 2013;76:403–6.
69. Yin Z, Gao J, Liu W, Huang C, Shuai X, Wang C, et al. Clinicopathological and prognostic analysis of primary gastrointestinal stromal tumor presenting with gastrointestinal bleeding: a 10-year retrospective study. *J Gastrointest Surg* 2017;21:792–800.
70. Maher ER, Neumann HP, Richard S. von Hippel-Lindau disease: a clinical and scientific review. *Eur J Hum Genet* 2011;19:617–23.
71. Eisenhofer G, Latke P, Herberg M, Siegert G, Qin N, Darr R, et al. Reference intervals for plasma free metanephrines with an age adjustment for normetanephrine for optimized laboratory testing of pheochromocytoma. *Ann Clin Biochem* 2013;50(Pt 1):62–9.
72. Franscini LC, Vazquez-Montes M, Budin T, Perera R, Dunand M, Grouzmann E, et al. Pediatric reference intervals for plasma free and total metanephrines established with a parametric approach: relevance to the diagnosis of neuroblastoma. *Pediatr Blood Cancer* 2015;62:587–93.
73. Criffin A, O'Shea P, FitzGerald R, O'Connor C, Torney W. Establishment of a paediatric age-related reference interval for the measurement of urinary total fractionated metanephrines. *Ann Clin Biochem* 2011;48(Pt 1):41–4.
74. Eisenhofer G, Goldstein DS, Walther MM, Friberg P, Lenders JW, Keiser HR, et al. Biochemical diagnosis of pheochromocytoma: how to distinguish true- from false positive test results. *J Clin Endocrinol Metab* 2003;88:2656–66.
75. McHenry CM, Hunter SJ, McCormick MT, Russell CF, Smye MC, Atkinson AB. Evaluation of the clonidine suppression test in the diagnosis of pheochromocytoma. *J Hum Hypertens* 2011;25:451–6.

Clinical Cancer Research

Von Hippel–Lindau and Hereditary Pheochromocytoma/Paraganglioma Syndromes: Clinical Features, Genetics, and Surveillance Recommendations in Childhood

Surya P. Rednam, Ayelet Erez, Harriet Druker, et al.

Clin Cancer Res 2017;23:e68-e75.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/23/12/e68>

Cited articles This article cites 70 articles, 8 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/23/12/e68.full#ref-list-1>

Citing articles This article has been cited by 1 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/23/12/e68.full#related-urls>

E-mail alerts Sign up to receive free email-alerts related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/23/12/e68>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.